



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 211/46, 211/40, A61K 31/40		A1	(11) International Publication Number: WO 95/24391
			(43) International Publication Date: 14 September 1995 (14.09.95)
(21) International Application Number: PCT/DK95/00111 (22) International Filing Date: 9 March 1995 (09.03.95) (30) Priority Data: 0277/94 9 March 1994 (09.03.94) DK (71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsvaerd (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): LUNDGREN, Karsten [DK/DK]; N. Jespersensvej 25, DK-2000 Frederiksberg (DK). KIRK, Ole [DK/DK]; Bisp Pedersvej 4, DK-2830 Virum (DK). (74) Common Representative: NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsvaerd (DK).		(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published With international search report.	


(54) Title: PIPERIDINES AND PYRROLIDINES

(57) Abstract

Compounds (I) and (II), the use of these compounds as medicament, the use of these medicaments in the treatment of diabetes and pharmaceutical compositions containing these compounds.



(I)



(II)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

PIPERIDINES AND PYRROLIDINES

Field of this invention

The present invention relates to novel compounds, the use of these compounds as medicament, the use of these medicaments
5 in the treatment of diabetes and pharmaceutical compositions containing these compounds.

Background of this invention

Diabetes is characterized by an impaired glucose metabolism manifesting itself among other things by an elevated blood
10 glucose level in the diabetic patients. Underlying defects lead to a classification of diabetes into two major groups: type 1 diabetes, or insulin demanding diabetes mellitus (IDDM), which arises when patients lack β -cells producing insulin in their pancreatic glands, and type 2 diabetes, or
15 non-insulin dependent diabetes mellitus (NIDDM), which occurs in patients with an impaired β -cell function besides a range of other abnormalities.

Type 1 diabetic patients are currently treated with insulin, while the majority of type 2 diabetic patients
20 are treated either with sulfonylureas that stimulate β -cell function or with agents that enhance the tissue sensitivity of the patients towards insulin or with insulin. Among the agents applied to enhance tissue sensitivity towards insulin, metformin is a representative example.

25 Even though sulfonylureas are widely used in the treatment of NIDDM, this therapy is, in most instances, not satisfactory: In a large number of NIDDM patients, sulfonylureas do not suffice to normalize blood sugar levels and the patients are, therefore, at high risk for acquiring diabetic
30 complications. Also, many patients gradually lose the ability to respond to treatment with sulfonylureas and are thus gradually forced into insulin treatment. This shift of

patients from oral hypoglycaemic agents to insulin therapy is usually ascribed to exhaustion of the β -cells in NIDDM patients.

In normals as well as in diabetics, the liver produces glucose in order to avoid hypoglycemia. This glucose production is derived either from the release of glucose from glycogen stores or from gluconeogenesis, which is a de novo intracellular synthesis of glucose. In type 2 diabetes, however, the regulation of hepatic glucose output is poorly controlled and is increased, and may be doubled after an overnight fast. Moreover, in these patients there exists a strong correlation between the increased fasting plasma glucose levels and the rate of hepatic glucose production (reviewed in R.A. De Fronzo: Diabetes 37 (1988), 667 - 687; A. Consoli: Diabetes Care 15 (1992), 430 - 441; and J.E. Gerich: Horm.Metab.Res. 26 (1992), 18 - 21). Similarly, hepatic glucose production will be increased in type 1 diabetes, if the disease is not properly controlled by insulin treatment.

Since existing forms of therapy of diabetes does not lead to sufficient glycaemic control and therefore are unsatisfying, there is a great demand for novel therapeutic approaches. Since the liver in diabetes is known to have an increased glucose production, compounds inhibiting this activity are highly desirable.

Recently, patents on inhibitors of the liver specific enzyme, glucose-6-phosphatase, which is necessary for the release of glucose from the liver, have been filed, for example German Offenlegungsschrift Nos. 4,202,183 and 4,202,184 and Japanese patent application No. 4-58565. All these known compounds are benzene derivatives.

International patent application having publication No. WO 92/16640 relates to di-, tri- and tetrasaccharides that are substrates or inhibitors of glycosyltransferase and glyco-

sidase enzymes. Some specific compounds mentioned therein are 2,3,4,5-tetrahydroxypiperidine, 3,4,5-trihydroxy-6-methylpiperidin and 3,4-dihydroxy-5-methylpiperidine.

International Patent Application No. WO 92/21657
5 relates to certain ω -deoxyazapyranoses, e.g. 3,4-dihydroxy-5-methylpiperidine mentioned in Claim 16 thereof. It is stated that these compounds have glucosidase inhibiting properties.

European patent application having publication No.
10 528,495 A1 relates to a class of azacyclic compounds, i.e. compounds comprising an azacyclic ring system substituted by arylmethoxy or an arylmethylthio moiety. These compounds may be useful as tachykinin antagonists.

European patent application having publication No.
15 375,651 A1 relates to 1,4-dideoxy-1,4-imino-L-allitol and derivatives thereof having glycosidase inhibitory activity.

European patent No. 947 B2 relates to derivatives of 3,4,5-trihydroxypiperidine having α -glucosidase inhibitory activity which can be used for the treatment of diabetes.

20 European patent No. 22,192 B1 relates to 1-alkadien-2,4-yl-2-hydroxymethyl-3,4,5-trihydroxypiperidines having maximal action on α -glucosidase hydrolases and lipid absorption which can be used for the treatment of diabetes.

European patent application having publication No.
25 481,950 A2 relates to 3-fluoro analogs of 2-hydroxymethyl-4,5-dihydroxypiperidines having glycosidase inhibition activity.

In J.Am.Chem.Soc. 113 (1991), 6678, 3,4-dihydroxypiperidine, 3,4-dihydroxy-5-methylpiperidine and 3,4-dihydroxy-6-
30 methylpiperidine are described as new compounds.

In J.Cell.Biochem., 1994, Suppl. 18D, 194 (published on 9 March 1994) some catalytic antibodies, e.g. (3R,4R,5R)-3,4-di-hydroxy-5-hydroxymethylpiperidine hydrochloride and methyl 6,7-dideoxy-7-((3R,4R,5R)-3,4-dihydroxy-5-hydroxy-
35 methylpiperidinyl)- α -D-glucoheptopyranoside hydrochloride, are described.

In Dansk Kemi 74 (1993), 31, it is mentioned that Ms. Jespersen is making a report (dissertation) concerning synthesis of (3R,4R,5R)-3,4-dihydroxy-5-hydroxymethylpiperidine and it is stated therein that, possibly, this report is
5 available. In this report, it is stated that this compound is a potential glycosidase inhibitor. Also in this report, 3-benzyloxy-4-hydroxy-5-hydroxymethylpiperidine is mentioned as an intermediate.

Moreover, scientifically it is well realized that in-
10 hibition of glycogen phosphorylase is a suitable target for the treatment of diabetes (Martin et al., 1991; Biochemistry 30: 10101-16; Oikonomakos et al., 1994; Eur. J. Drug Metab. Pharmacokin. 3: 185-92). These groups have used glucose ana-
logs.

15 One object of the present invention is to furnish compounds which can be used as medicaments.

A further object of this invention is to furnish compounds which can effectively be used in the treatment of diabetes.

20 A still further object of this invention is to furnish compounds which can effectively be used as inhibitors of glucose production from the liver.

Brief description of this invention

Surprisingly, it has been found that compounds of the
25 general formulas I and II stated in the claims, below, have interesting pharmacological properties. For example, the compounds of this invention can be used in the treatment of diabetes. Especially, the compounds of this invention are active as inhibitors of glucose production from the liver.
30 Consequently, the compounds of this invention can be used for the treatment of the increased plasma glucose levels in diabetics.

Detailed description of this invention

Hereinafter, the term alkyl, when used alone or in combination with another moiety, is a straight or branched alkyl group which preferably contains not more than 8 carbon atoms, more preferred not more than 4 carbon atoms. Especially preferred alkyl groups are methyl, ethyl, propyl and isopropyl.

Preferably, hydroxyalkyl is hydroxymethyl or hydroxyethyl. Preferably, alkoxy is methoxy or ethoxy. Preferably, halogen is chloro, bromo or fluoro. Preferably, alkylamino is methylamino. Preferably, acylamino is alkylcarbonylamino, more preferred acetylamino. Preferably, dialkylamino is N,N-dimethylamino. Preferably, carboxyalkyl is methoxycarbonyl. Preferably, alkylthio is methylthio or ethylthio. Preferably, alkenyl is ethenyl or propenyl. Preferably, alkylphenyl is o-, m- or p-tolyl.

When the term trialkyl ammonium ion is used herein without stating which ion it is connected to, it is to be understood that it is connected to an ion, preferably the residue of an acid from which, for example, a hydrogen atom has been removed. Similar considerations apply for the terms quaternary ammonium base ion and N,N,N-trialkylammonium ion.

The term monosaccharide moiety designates a monosaccharide from which a hydrogen atom has been removed. Examples of preferred monosaccharides usable to form such moieties (R^1) are glucose, fructose, galactose, mannose and methyl 6,7-dideoxy-D-glucopyranoside.

Examples of preferred compounds of formula I and II are (3R,4R,5R)-3,4-dihydroxy-5-hydroxymethylpiperidine, 3,4-dihydroxy-5-hydroxymethylpiperidine, 3-benzyloxy-4-hydroxy-5-hydroxymethylpiperidine, 3,4-dihydroxy-5-hydroxymethylpiperidine N-(7-(methyl 6,7-dideoxy-D-glucopyranoside)), 3,4-dihydroxy-5-methylpiperidine, 3,4-dihydroxy-5-

ethylpiperidine, 3,4-dihydroxy-5-propylpiperidine, 3,4-dihydroxy-5-isopropylpiperidine, 3,4-dihydroxy-5-phenylpiperidine, 3,4-dihydroxy-5-hydroxyethylpiperidine, 3,4-dihydroxy-5-fluoromethylpiperidine, 3,4-dihydroxy-5-chloromethylpiperidine, 3-hydroxy-5-hydroxymethylpiperidine, 3-hydroxy-4-fluoro-5-hydroxymethylpiperidine, 3-hydroxy-4-chloro-5-hydroxymethylpiperidine, 3-fluoro-4-hydroxy-5-hydroxymethylpiperidine, 3-chloro-4-hydroxy-5-hydroxymethylpiperidine, 4-hydroxy-5-hydroxymethylpiperidine, N-methyl-3,4-dihydroxy-5-hydroxymethylpiperidine, N,N-dimethyl-3,4-dihydroxy-5-hydroxymethylpiperidinium chloride, (3R,4R)-3-hydroxy-4-hydroxymethylpyrrolidine, 3-hydroxy-4-hydroxymethylpyrrolidine, 3-benzyloxy-4-hydroxymethylpyrrolidine, 3-hydroxy-4-hydroxymethylpyrrolidine N-(7-(methyl 6,7-dideoxy-D-glucopyranoside)), 3-hydroxy-4-methylpyrrolidine, 3-hydroxy-4-ethylpyrrolidine, 3-hydroxy-4-propylpyrrolidine, 3-hydroxy-4-isopropylpyrrolidine, 3-hydroxy-4-phenylpyrrolidine, 3-hydroxy-4-hydroxyethylpyrrolidine, 3-hydroxy-4-fluoromethylpyrrolidine, 3-hydroxy-4-chloromethylpyrrolidine, 3-hydroxy-4-hydroxymethylpyrrolidine, 3-fluoro-4-hydroxymethylpyrrolidine, 3-chloro-4-hydroxymethylpyrrolidine, N-methyl-3-hydroxy-4-hydroxymethylpyrrolidine and N,N-dimethyl-3-hydroxy-4-hydroxymethylpyrrolidinium chloride.

The compounds of formulas I and II may be presented as a mixture of enantiomers which, if desired, may be resolved into the individual pure enantiomers. This resolution may conveniently be performed by fractional crystallization from various solvents, of the salts of compounds of the formula I or II with optical active acids or by other methods known *per se*, for example, chiral column chromatography. This invention includes all isomers, whether resolved or mixtures thereof.

Examples of pharmaceutically acceptable salts are acid addition salts with non-toxic acids, either inorganic acids

such as hydrochloric acid, sulphuric acid and phosphoric acid, or organic acids such as formic acid, acetic acid, propionic acid, succinic acid, gluconic acid, lactic acid, citric acid, ascorbic acid, benzoic acid, embonic acid, methanesulphonic acid and malonic acid.

The compounds of formulas I and II are prepared by methods known per se by the skilled art worker, for example as described in the following. One synthetic strategy can be to employ a chiral Cerny epoxide (Tanka, T. and Cerny, M. Collection Czechoslov. Chem. Commun: 36 (1971), 2216), which is opened by a reagent to introduce a hydroxymethyl group (as illustrated in Example 1, below) or another alkyl or substituted alkyl substituent. After hydrolysis of the anhydro bond, the pentodialdose could be obtained by oxidative cleavage of the carbon chain. As illustrated in Example 1, (3R,4R,5R)-3,4-dihydroxy-5-hydroxymethylpiperidine was prepared by reductive amination of the pentodialdose. Similarly, other compounds of the general formula I or II can be prepared by the above strategy. A variety of functional groups can be introduced in the compounds prepared as outlined above by methods well known to those skilled in the art.

Pharmaceutical Compositions

This invention further provides pharmaceutical compositions which comprise at least one compound of formula I or II or a pharmaceutically acceptable salt thereof in connection with a pharmaceutically acceptable carrier. Such compositions may be in the form of powders, solutions, or suspensions, which may or may not be divided in unit dosage form or in the form of capsules or tablets.

The pharmaceutical compositions of this invention may comprise carriers, diluents, absorption enhancers, tablet disintegrating agents and other ingredients which are conventionally used in the art. The powders and tablets preferably

contain from 5 to 99%, more preferred from 10 to 90 % of the active ingredient. Examples of solid carriers are magnesium carbonate, magnesium stearate, dextrin, lactose, sugar, talc, gelatin, pectin, tragacanth, methyl cellulose, sodium 5 carboxymethyl cellulose, low melting waxes and cocoa butter.

Liquid compositions include sterile solutions, suspensions and emulsions suitable for parenteral injection.

The route of administration of the compositions containing a compound of formula I or II may be any route which 10 effectively transports the active compound to its site of action, the oral or nasal route being preferred.

The regimen for any patient to be treated with the compositions according to the present invention should be determined by those skilled in the art. The daily dose to be 15 administered in therapy can be determined by a physician and will depend on the particular compound employed, on the route of administration and on the age and the condition of the patient. A convenient daily dosage can be less than about 1 g, preferably in the range around 10 - 200 mg.

20 The present invention is further illustrated by the following examples which, however, are not to be construed as limiting the scope of protection. The features disclosed in the foregoing description and in the following examples may, both separately and in any combination thereof, be 25 material for realising the invention in diverse forms thereof.

Example 1

Synthesis of (3R,4R,5R)-3,4-dihydroxy-5-hydroxymethylpiperidine.

30 The title compound was synthesized by a 6 step procedure starting from 1,6:2,3-dianhydro-4-O-benzyl- β -D-mannopyranose as outlined below:

1,6-Anhydro-4-O-benzyl-2-deoxy-2-C-vinyl- β -D-glucopyranose:

To a stirred solution of 1,6:2,3-dianhydro-4-O-benzyl- β -D-mannopyranose (prepared as outlined by Tanka, T. and Cerny, M. Collection Czechoslov. Chem. Commun.: 36 (1971), 2216) 5 (5.0 g, 21.3 mmol) in dry tetrahydrofuran (hereinafter designated THF) (25 ml), was added a solution of 1.77 M vinylmagnesiumbromide in THF (120.5 ml, 213 mmol). The mixture was gently refluxed for 3 h at 60°C. After cooling to room temperature, a 2 M NH_4Cl aqueous solution (650 ml, pH 10 value: 8) was slowly added. The aqueous layer was extracted with ethyl acetate (2 x 250 ml). The combined organic layers were dried (MgSO_4), filtered and concentrated. The crude product (4.87 g, 87%) was purified by flash chromatography using ethyl acetate/pentane (1:2) as eluent, to give 15 1,6-anhydro-4-O-benzyl-2-deoxy-2-C-vinyl- β -D-glucopyranose as a white crystalline compound in 75.5% (4.23 g) yield. After recrystallization in ethanol, the melting point was 88 - 91°C. ^{13}C -NMR (50 MHz, CDCl_3): δ 135.5 (CH=); 137.7, 128.3, 127.6, 127.5 (phenyl, hereinafter designated Ph); 20 117.2 ($\text{CH}_2=$); 103.5 (C-1); 78.8 (C-4); 74.6 (C-3); 71.3, 70.4 (C-5, OCH_2Ph); 65.4 (C-6); 51.8 ppm (C-2). ^1H NMR (200 MHz, CDCl_3): d 7.35 (d, 5H, Ph); 5.99 (m, 1H, CH=CH_2); 5.25 and 5.17 (s, 1H and d, 1H, $\text{CH}_2=$); 4.63 (dd, 3H, OCH_2Ph and H-5); 4.09 and 3.75 (d, 1H og m, 2H, H-3, H-6 og H-6'); 3.43 25 (s, 1H, H-4); 2.80 (broad s, 1H, OH); 2.45 (d, 1H, H-2). Analytical calculation for $\text{C}_{15}\text{H}_{18}\text{O}_4$: C: 68.69; H: 6.92. Found: C: 68.58; H: 6.94.

1,6-Anhydro-4-O-benzyl-2-deoxy-2-C-hydroxymethyl- β -D-glucopyranose:

30 A stream of ozone (0.36 mmol/minute) was passed through a solution of 1,6-anhydro-4-O-benzyl-2-deoxy-2-C-vinyl- β -D-glucopyranose, (3.36 g, 12.8 mmol) in ethanol (100 ml) for 43.5 minutes. The solution of the ozonide was transferred to

a three necked round bottomed flash equipped with thermometer and a dropping funnel containing a solution of NaBH_4 (3.87 g, 102 mmol) in ethanol/water (1:1; 35 ml). The solution of NaBH_4 was added dropwise in such a manner to keep the temperature below 20°C (cooling on ice bath). After stirring for 45 minutes, Amberlite IR 120, H^+ (100 ml) was added. After stirring for another 30 minutes, the ion exchange resin was filtered off and rinsed with water. The solution was evaporated and co-concentrated with methanol (3 x 80 ml). The residue was a colourless sirup in 96% (3.28 g) yield. This sirup was flash chromatographed using ethyl acetate/pentane (2:1) and ethyl acetate as eluent to give the title compound in a 63% (2.14 g) yield as a crystalline compound. In some runs, the crude sirup crystallised and was recrystallised from chloroform: melting point $111 - 113^\circ\text{C}$. ^{13}C -NMR (50 MHz, D_2O): δ 139, 130.1, 129.8 (Ph); 102.0 (C-1); 80.5 (C-4); 76.1 (C-3); 72.9 (OCH_2Ph); 67.4 (C-5); 66.2 (C-6); 61.7 (C-2'); 49.0 ppm (C-2). Analytical calculation for $\text{C}_{14}\text{H}_{18}\text{O}_5$: C: 63.15; H: 6.81. Found: C: 63.14; H: 6.79.

4-O-Benzyl-2-deoxy-2-C-hydroxymethyl-D-glucopyranose:

1,6-Anhydro-4-O-benzyl-2-deoxy-2-C-hydroxymethyl- β -D-glucopyranose (1.59 g, 6.0 mmol) was dissolved in 1 M sulfuric acid (30 ml) by heating to reflux. The anhydride was refluxed for 1 hour. After cooling to room temperature, the solution was poured through a column containing Amberlite[®] IR 67, OH^- (120 ml). The column was rinsed with water followed by methanol (400 ml). The eluated liquid was concentrated. The residue was a colourless sirup obtained in 98% (1.67 g) yield. Flash chromatography using ethyl acetate and ethyl acetate/methanol (10:1) as eluent gave the product in 76% (1.29 g) yield. One of the anomers (β) could be obtained by crystallization in ethyl acetate in 27% yield as a white crystalline compound. Melting point: $102 - 105^\circ\text{C}$. ^{13}C -NMR (50 MHz, D_2O): δ 136.8, 128.3, 128.0 (Ph); 94.1

(C-1, β); 91.4 (C-1, α); 78.6 (C-4); 74.4 (C-3); 70.3 (OCH₂Ph, C-6 β); 69.5 (C-6, α); 60.3 (C-5); 59.1 (C-2', α); 56.6 (C-2', β); 47.7 ppm (C-2, α). ¹H-NMR (200 MHz, D₂O); 5.13 (d, 1H, H-1 (α), $J_{1,2}$ = 3.5 Hz); 4.48 (d, 1H, H-1 (β), $J_{1,2}$ = 10 Hz). Analytical calculation for C₁₄H₂₀O₆ x 0.3 H₂O: C: 58.04; H: 7.17. Found: C: 58.07; H: 7.19.

4-O-Benzyl-2-deoxy-2-C-hydroxymethyl-D-xylo-pentodialdose:

To a solution of 4-O-benzyl-2-deoxy-2-C-hydroxymethyl-D-gluco-pyranose (1.41 g, 5.0 mmol) in methanol (15 ml), a
10 solution of sodium periodate (5.35 g, 25.0 mmol) in water (50 ml) was added dropwise over 15 minutes. Further, methanol (25 ml) was added. The mixture was stirred at 45°C for 3 hours. The precipitated iodate was filtered off, and the mixture was concentrated. By resolution of the residue
15 in ethyl acetate/ethanol (1:1; 80 ml), more iodate was precipitated and filtered off. The mother liquor was concentrated and the residue (2.07 g) was flash chromatographed using ethyl acetate as eluent. The purified product was obtained as a yellow sirup in 87% (1.09 g) yield and 9% of
20 unreacted starting material (0.13 g) was isolated. The purified product was used immediately in the next reaction.

(3R,4R,5R)-3-Benzyl-4-hydroxy-5-hydroxymethylpiperidine:

To a solution of 4-O-benzyl-2-deoxy-2-C-hydroxymethyl-D-xylo-pentodialdose (1.77 g) in ethanol (40 ml), 0.29 M NH₃
25 in ethanol (162 ml) and 5% palladium on charcoal (300 mg) was added. The mixture was hydrogenated at 500 Psi at 20°C for 15 hours. The reaction mixture was filtrated and concentrated. The residue (1.85 g) was flash chromatographed using ethanol/NH₄OH (25% aqueous)/triethylamine (122:2:1) as
30 eluent giving the product as a colourless sirup (getting coloured after storing) in 78% yield. ¹³C-NMR (62.9 MHz, D₂O): δ 138.1, 128.1, 127.3 (Ph); 80.6 (C-3), 74.9 (C-4);

71.8 (OCH₂Ph); 62.5 (C-5'); 48.0 (C-2); 46.8 (C-6); 45.0 ppm (C-5). 1H-NMR (500 MHz, D₂O): δ 7.3 (s, 5H, Ph); 4.65, 4.51 (2 d, 2H, OCH₂Ph, $J_{\text{gem}} = 12$ Hz); 3.66 (dd, 1H, H-5a', $J_{5a',5b'} = 10$ Hz, $J_{5a',5} = 5.5$); 3.57 (dd, 1H, H-5'b, $J_{5a',5b'} = 10$ Hz, $J_{5b',5} = 4.5$); 3.41 (dd, 1H, H-4, $J_{3,4} = 11$ Hz, $J_{4,5} = 9$ Hz); 3.3 (broad, N-H); 3.27 (dd, 1H, H-3, $J_{3,2zx} = 11$ Hz, $J_{3,2eq} = 4$ Hz); 3.22 (dd, 1H, H-2eq, $J_{2eq,2ax} = 11$, $J_{2eq,3} = 4$ Hz); 2.97 (dd, 1H, H-6 eq, $J_{6eq,6ax} = 12$ Hz, $J_{6eq,5} = 4$ Hz); 2.38 (t, 1H, H-2ax, $J_{2ax,2eq} = J_{2ax,3} = 11$ Hz); 2.31 (t, 1H, H-6ax, $J_{6ax,6eq} = J_{6ax,5} = 12$ Hz); 1.80 ppm (m, 1H, H-5).

(3R,4R,5R)-3,4-Dihydroxy-5-hydroxymethylpiperidine hydrochloride:

(3R,4R,5R)-3-Benzyloxy-4-hydroxy-5-hydroxymethylpiperidine (0.527 g, 2.2. mmol) was dissolved in 0.5 M HCl (5.3 ml) and ethanol (50 ml) and 5% palladium charcoal (300 mg) was added. The mixture was hydrogenated at 101 kPa and 20°C for 18 hours. The reaction mixture was filtrated and concentrated to give the product in 93% (0.375 g) yield.

13C-NMR (50 MHz, D₂O): δ 70.7 og 68.1 (C-3 og C-4); 58.6 (C-5'); 46.2 og 44.4 (C-2 og C-6); 40.6 (C-5). 1H-NMR (500 MHz, D₂O, Ph < 1, ref. 4.63 ppm): δ 3.72 (dd, 1H, H-5b', $J_{5a',5b'} = 11.5$, $J_{5,5b'} = 3.3$ Hz); 3.67 (ddd, 1H, H-3, $J_{3,2ax} = 11.2$, $J_{3,4} = 8.9$, $J_{3,2eq} = 4.9$ Hz); 3.64 (dd, 1H, H-5a', $J_{5a',5b'} = 11.5$, $J_{5a',5} = 6.2$ Hz); 3.43 (ddd, 1H, H-2eq, $J_{2eq,2ax} = 12.7$, $J_{2eq,3} = 4.9$, $J_{2eq,6eq} = 2.0$); 3.42 (dd, 1H, H-4, $J_{4,5} = 10.5$, $J_{4,3} = 8.9$ (Hz); 3.41 (ddd, 1H, H-6eq, $J_{6eq,6ax} = 13.4$, $J_{6eq,5} = 3.8$, $J_{6eq,2eq} = 2.0$ Hz); 2.87, 12.7, $J_{2ax,3} = 11.2$ Hz); 1.86 ppm (dddd, 1H, H-5). If necessary, the piperidine could be chromatographed using ethanol/NH₄OH (25% aqueous) (10:1) to give the free piperidine. Analytical calculation for C₆H₁₃NO₃: C: 48.97; H: 8.90; N: 9.52. Found: C: 48.46; H: 9.33; N: 9.17.

Example 2

Experimental protocol and results.

Rat hepatocytes were isolated using a standard two step collagenase technique, and cultured onto collagen coated culture dishes for 72 hours in medium 199 with the addition of dexamethazone (0.1 μ M); penicillin/Streptomycin ((100 u/100 μ g)/ml) and insulin (1 nM). During the last 24 hours, the hepatocytes were cultured in the presence of high levels of insulin (5 nM) and glucose (15 mM), which result in the incorporation of glucose into glycogen. Therefore, at the time of the experiment, the cells mimic livers from fed animals.

Experiments were initiated after 72 hours of culture by 2 times wash of cells and addition of a 20 mM HEPES experimental buffer including balanced salts, but without glucose. The test compound was added simultaneously with the experimental buffer. To some cultures, glucagon (0.5 nM) was added after 10 minutes in order to stimulate glucose production from liver cells. The glucose released into the media, reflecting the glucose production of the liver cells, was measured 70 minutes after the start of the experiment and standardized to cellular DNA content.

Phosphorylase was either purchased from Sigma or extracted from rat livers according to Stalmans et. al. (Eur. J. Biochem. 49 (1974), 415). The activity of phosphorylase was determined as described by Bergmeyer (1983; in : Meth. of Enzymatic Analysis, 2 pp 293 - 295, Weinheim, (ed.) Verlag Chemie).

The activity of the glycogen debranching enzyme, α -1,6-glucosidase was determined as described by Brown and Brown (1966; in : Meth. in Enzymology, 8 : 515 - 524, Neufeld and Ginsburg (Eds.) Academic Press).

Table 1 below shows the results obtained with the compound of example 1 on basal and glucagon stimulated glycogenolysis. The effects are compared to those exerted by the model debranching enzyme inhibitor 1-deoxynojirimycin 5 (hereinafter designated dNOJ).

10

Table 1. Effects of Example 1 on baseline and glucagon stimulated glucose production from cultured liver cells. Values are expressed relative to the basal glucose production. Results obtained with the model α -1,6 glucosidase inhibitor 1-deoxynojirimycin are shown for comparison.

	<u>Glucose production</u>	
	Without glucagon	With glucagon (0.5nM)
No addition:	100 %	283%
Example 1 (32 μ M):	17 %	32%
1-deoxynojirimycin (50 μ M):	89 %	237%

15 The results clearly demonstrate the ability of the compound of example 1 to inhibit basal and glucagon stimulated hepatocyte glucose production, while dNOJ only exerted marginal effects.

20 Table 2 compares the potency of the compound of example 1 with the potency of dNOJ on various cellular and enzymatic activities.

5 **Table 2. Effects of compound of example 1 compared to effects of 1-deoxynojirimycin on different cellular and enzymatic activities. The results are the concentrations of the compounds resulting in half maximal activity (I.C.₅₀).**

	I.C. ₅₀ values (μmoles/L)	
	Example 1	1-deoxynojirimycin
Liver cell glucose production:		
Basal:	1.3	> 100
Glucagon stimulated:	1.6	> 100
10 Phosphorylase:	1.12	> 200
liver α-1,6-glucosidase	77	2.4
Yeast α-1,6-glucosidase*	85.9	25.4

*From: Jespersen et al. (Angew. Chem. Int. Ed. Engl. 33 (1994), 1778)

15 It is apparent from the presented data in table 2 that the compound of example 1 is a potent inhibitor of liver cell glucose production. Moreover, it is also demonstrated that phosphorylase is inhibited by this compound in similar low concentrations. In contrast, the compound of example 1 is
20 only a very weak inhibitor of α-1,6-glucosidase either from liver or from yeast.

Table 2 also demonstrates that the potent model inhibitor of liver α-1,6-glucosidase was unable to inhibit either liver cell glucose production or phosphorylase. These data are in
25 agreement with data obtained by Bollen and Stalmans (Eur. J. Biochem. 181 (1980), 775), who also concluded that α-1,6-glucosidase inhibition is an insufficient principle for inhibition of liver cell glucose production.

In conclusion, the data demonstrates that inhibition of
30 liver cell glucose production by the compounds of formula I

and II according to this invention, as exemplified by the compound of example 1, is mediated by inhibition of phosphorylase. Consequently, the compounds of formula I and II can be used to inhibit both the base line and glucagon stimulated glucose production from liver cells.

Example 3

Tablets

Tablets which are suitable for oral administration and which contain the below-mentioned components are produced in a manner known per se granulating the active and the auxiliary substances and making them into tablets.

Each tablet contains 50 mg of the compound of formula I or II, e.g. the compound of Example 1, 100 mg of lactose, 30 mg of corn starch, 3 mg of talc powder, 3 mg of colloidal silicon dioxide and 2 mg of magnesium stearate.

Example 4


Capsules

Capsules which are suitable for oral administration contain the below-mentioned components are produced in a manner known per se mixing the active substances with the auxiliary substances and putting them into gelatine capsules.

Each capsule contains 50 mg of the compound of formula I or II, e.g. the compound of example 1, 100 mg of lactose, 30 mg of corn starch, 3 mg of talc powder, 3 mg of colloidal silicon dioxide and 2 mg of magnesium stearate.

Claims:

1. Compounds of the general formulas I and II according to Claim 3 or 7, respectively.
2. Novel compounds according to Claim 1.
- 5 3. Compounds, according to Claim 1 or 2, having the general formula I



wherein R^1 represents hydrogen, a monosaccharide moiety or alkyl which optionally is substituted by one or more of the following groups: hydroxy, hydroxyalkyl, halogen, amino, 10 alkylamino, dialkylamino, a trialkylammonium ion, nitro, formyl, carboxy, carboxyalkyl, alkylthio, alkenyl, phenyl and alkylphenyl, or R^1 together with the adjacent nitrogen atom from the piperidine nucleus represents a quaternary ammonium base ion residue containing two alkyl groups which 15 optionally are substituted by one or more of the following groups: hydroxy, hydroxyalkyl, halogen, amino, alkylamino, dialkylamino, a trialkylammonium ion, nitro, formyl, carboxy, carboxyalkyl, alkylthio, alkenyl, phenyl and alkylphenyl; R^2 , R^3 , R^4 and R^6 , which are the same or different, 20 independent of each other represents hydrogen, hydroxy, hydroxyalkyl, halogen, amino, alkylamino, acylamino, N,N-


dialkylamino, a N,N,N-trialkylammonium ion, nitro, formyl, carboxy, benzoxy, mercapto, alkylthio, alkenyl, phenyl and alkylphenyl; and R⁵ represents phenyl or methyl which optionally is substituted by one or more of the following groups: alkyl, hydroxy, hydroxyalkyl, halogen, amino, alkyl-
5 amino, dialkylamino, trialkylammonium, nitro, formyl, carboxy, carboxyalkyl, alkylthio, alkenyl, phenyl and alkylphenyl; with the proviso that said compound contains at least 2 free or protected hydroxy groups, preferably 2 free
10 hydroxy groups; and salts thereof.

4. Compounds, according to anyone of the previous claims with the proviso that it is different from 3,4-dihydroxy-5-methylpiperidine (3R,4R,5R)-3,4-dihydroxy-5-hydroxymethylpiperidine, and 3-benzyloxy-4-hydroxy-5-hydroxymethylpiper-
15 idine.

5. Compounds, according to any one of the preceding claims, containing at least 3 free or protected hydroxy groups, preferably 3 free hydroxy groups.

6. Compounds, according to the previous claim, wherein the
20 optionally protected hydroxy groups are the substituents R³, R⁴ and R⁵ or are present in said substituents.

7. Compounds, according to Claim 1 or 2, having the general formula II



wherein R^1 represents hydrogen, a monosaccharide moiety or alkyl which optionally is substituted by one or more of the following groups: hydroxy, hydroxyalkyl, halogen, amino, alkylamino, dialkylamino, a trialkylammonium ion, nitro, formyl, carboxy, carboxyalkyl, alkylthio, alkenyl, phenyl and alkylphenyl, or R^1 together with the adjacent nitrogen atom from the pyrrolidine nucleus represents a quaternary ammonium base ion residue containing two alkyl groups which optionally are substituted by one or more of the following groups: hydroxy, hydroxyalkyl, halogen, amino, alkylamino, dialkylamino, a trialkylammonium ion, nitro, formyl, carboxy, carboxyalkyl, alkylthio, alkenyl, phenyl and alkylphenyl; R^2 , R^3 and R^5 , which are the same or different, independent of each other represents hydrogen, hydroxy, hydroxyalkyl, halogen, amino, alkylamino, N,N-dialkylamino, a N,N,N-trialkylammonium ion, nitro, formyl, carboxy, benzoy, mercapto, alkylthio, alkenyl, phenyl and alkylphenyl; and R^4 represents carboxy, phenyl or methyl which optionally is substituted by one or more of the following groups: alkyl, hydroxy, hydroxyalkyl, halogen, amino, alkylamino, dialkylamino, a trialkylammonium ion, nitro, formyl, carboxy, carboxyalkyl, alkylthio, alkenyl, phenyl and alkylphenyl; with the proviso that said compound contains at

least 2 free or protected hydroxy groups, preferably 2 free hydroxy groups; and salts thereof.

8. Compounds, according to the previous claim containing at least 3 free or protected hydroxy groups, preferably 3
5 free hydroxy groups.

9. Compounds, according to Claim 7, wherein the optionally protected hydroxy groups are substituents R^3 and R^4 or are present in said substituents.

10. Compounds, according to any one of the Claims 3 through
10 6, wherein R^1 is hydrogen, a monosaccharide moiety or alkyl, preferably methyl, or R^1 together with the adjacent nitrogen atom is a dialkylammonium ion, preferably a dimethylammonium ion.

11. Compounds, according to any one of the Claims 3 through
15 6 and 10, wherein R^2 is hydrogen.

12. Compounds, according to any one of the Claims 3 through 6, 10 and 11, wherein R^3 is hydroxy, halogen, preferably fluoro or chloro, or benzyloxy.

13. Compounds, according to any one of the Claims 3 through
20 6 and 10 through 12, wherein R^4 is hydroxy or halogen, preferably fluoro or chloro.

14. Compounds, according to any one of the Claims 3 through 6 and 10 through 13, wherein R^5 is hydroxymethyl, hydroxy-ethyl, methyl, ethyl, propyl, isopropyl, fluoromethyl,
25 chloromethyl or phenyl.

15. Compounds, according to any one of the Claims 3 through 6 and 10 through 14, wherein R^6 is hydrogen.

16. Compounds, according to any one of the Claims 7 through 9, wherein R^1 is hydrogen, a monosaccharide moiety or alkyl, preferably methyl, or R^1 together with the adjacent nitrogen atom is a dialkylammonium ion, preferably a dimethylammonium ion.
5

17. Compounds, according to any one of the Claims 7 through 9 and 16, wherein R^2 is hydrogen.

18. Compounds, according to any one of the Claims 7 through 9, 16 and 17, wherein R^3 is hydroxy, halogen, preferably fluoro or chloro, or benzyloxy.
10

19. Compounds, according to any one of the Claims 7 through 9 and 16 through 18, wherein R^4 is hydroxymethyl, hydroxyethyl, methyl, ethyl, propyl, isopropyl, fluoromethyl, chloromethyl or phenyl.

15 20. Compounds, according to any one of the Claims 7 through 9 and 16 through 19, wherein R^5 is hydrogen.

21. The compound, according to Claim 1, which is 3,4-dihydroxy-5-hydroxymethylpiperidine and any of the optical isomers thereof.

20 22. The compound, according to the previous claim, which is (3R,4R,5R)-3,4-dihydroxy-5-hydroxymethylpiperidine.

23. The use of a compound of formula I or II according to any one of the preceding claims or anyone of the specific compounds mentioned herein and covered by formula I or II as
25 medicament.

24. The use of a compound of formula I or II according to anyone of Claims 1 through 22 or anyone of the specific com-

pounds mentioned herein and covered by formula I or II as a liver glycogen phosphorylase inhibitory agent.

25. The use of a compound of formula I or II according to any one of the Claims 1 through 22 or anyone of the specific
5 compounds mentioned herein and covered by formula I or II for the manufacture of a pharmaceutical composition for the treatment of diabetes.

26. The use of a compound of formula I or II according to anyone of the claims 1 through 22 or anyone of the specific
10 compounds mentioned herein and covered by formula I or II for the manufacture of a pharmaceutical composition inhibiting the glucose production from the liver.

27. A pharmaceutical composition containing a compound of formula I or II according to any one of the Claims 1 through
15 22 or anyone of the specific compounds mentioned herein and covered by formula I or II in connection with a pharmaceutically acceptable carrier.

28. A method of treating diabetes which method comprises administering an effective amount of a compound of formula I
20 or II according to any one of the Claims 1 through 22 or anyone of the specific compounds mentioned herein and covered by formula I or II to a patient in need of such a treatment.

29. A method of inhibiting the liver glucose production
25 from the liver which method comprises administering an effective amount of a compound of formula I or II according to anyone one the claims 1 through 22 or anyone of the specific compounds mentioned herein and covered by formula I or II to a patient in need of such a treatment.

30. Any novel feature or combination of features described herein.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 95/00111

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07D 211/46, C07D 211/40, A61K 31/40

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS-ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO, A1, 9221657 (THE SCRIPPS RESEARCH INSTITUTE), 10 December 1992 (10.12.92), see especially pages 38-34 --	1-19,25-27
X	EP, A1, 0000947 (BAYER AKTIENGESSELLSCHAFT), 7 March 1979 (07.03.79) --	1-6,10-14, 25-27
X	EP, A2, 0481950 (MONSANTO COMPANY), 22 April 1992 (22.04.92) --	1-6,10-14, 25-27
X	EP, A1, 0375651 (MONSANTO COMPANY), 27 June 1990 (27.06.90) --	1-2,7-9, 16-20,25-27

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

20 June 1995

Date of mailing of the international search report

26 -06- 1995

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Göran Karlsson
Telephone No. +46 8 782 25 00

2
INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 95/00111

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>EP, A1, 0022192 (BAYER AG), 14 January 1981 (14.01.81)</p> <p style="text-align: center;">-- -----</p>	<p>1-6, 11-14, 25-27</p>

INTERNATIONAL SEARCH REPORT

Information on patent family members

03/05/95

International application No.

PCT/DK 95/00111

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A1- 9221657	10/12/92	AU-A- 2145992 US-A- 5276120	08/01/93 04/01/94
EP-A1- 0000947	07/03/79	SE-T3- 0000947 AT-B- 373239 AT-B- 376420 AT-B- 376421 AU-B- 520686 AU-A- 3921478 AU-A- 3930478 CA-A- 1123437 DE-A- 2738717 JP-C- 1442470 JP-A- 54046786 JP-B- 62031703 SU-A, A- 917697 US-A- 4260622 US-A- 4639436 DE-A- 2758025 SU-A- 1014470 SU-A- 1087074	27/12/83 26/11/84 26/11/84 18/02/82 28/02/80 06/03/80 11/05/82 08/03/79 08/06/88 12/04/79 09/07/87 30/03/82 07/04/81 27/01/87 12/07/79 23/04/83 15/04/84
EP-A2- 0481950	22/04/92	JP-A- 4257562 US-A- 5128347 US-A- 5273981	11/09/92 07/07/92 28/12/93
EP-A1- 0375651	27/06/90	SE-T3- 0375651 AT-T- 112558 CA-A- 2006269 DE-D, T- 68918681 ES-T- 2064487 JP-A- 2212470 US-A- 4894388	15/10/94 22/06/90 30/03/95 01/02/95 23/08/90 16/01/90
EP-A1- 0022192	14/01/81	SE-T3- 0022192 AT-T- 3421 AU-A- 5933080 CA-A- 1138879 DE-A- 2925943 JP-C- 1303453 JP-A- 56007763 JP-B- 60026388 US-A- 4312872	15/06/83 08/01/81 04/01/83 29/01/81 28/02/86 27/01/81 24/06/85 26/01/82